P158 (continued)

Guidelines. However, fresh vegetables were slightly more common in households than fresh fruits. Canned, frozen, and packaged fruits, vegetables, and legumes, were more prevalent than fresh forms. The major protein source was red meat or breakfast meat. Whole grain was less common than white, refined grain products.

Conclusions and Implications: Mothers and children in substance abuse recovery programs could benefit from nutrition education and improved access to healthier food items. Further research focusing on shifts in addition and food choice would be beneficial to guide nutrition interventions and better understand the process of recovery among mothers in substance abuse recovery facilities.

Funding: East Carolina University Office of Engagement, Innovation, and Economic Development

P159 The Efficacy of Using Facebook as a Channel on the School Weight Control Program

Yaolin Weng, PhD, ylweng@pu.edu.tw, Providence University, 200 Zhongqi Road, Shalu Township, Taichung City, 43301, Taiwan; T. L. Kang, MS

Objective: The amount of time that young adults spent online has been doubled, and most of them were using Facebook. Thus, the purpose of this study is to discuss the efficacy of using Facebook social networking sites as an intervention channel on the school weight control program among the university students.

Study Design, Setting, Participants, and Intervention: Fifty students with BMI 24kg/m² were recruited and randomly assigned into two groups for eight weeks, one group accepted regular nutrition education and the other group participated the Facebook intervention. Only the Facebook intervention group accepted nutrition newsletters once a week through Facebook network and participated one hour on-line discussion section every day regarding to weight loss tips and questions.

Outcome, Measures and Analysis: Both groups were pre and post-tested with the self-designed survey and the anthropometric measurements. KAP scores and anthropometric measurements were compared by pair-t test for both groups. Login times and discussion contents among the Facebook group were recorded for qualitative analyses.

Results: The mean weight loss in the Facebook group was 2.8 kg ($p < 0.001$), and the body fat was decreased 3.2% ($p < 0.001$). Moreover, the nutrition knowledge related to weight-control was increased 0.7 points ($p < 0.001$), the weight-control attitude was increased 1.4 points ($p < 0.001$), and the weight-control behavior was increased 4.4 points ($p < 0.001$).

Conclusions and Implications: In summary, both regular and Facebook intervention could increase nutrition knowledge, improve the weight-control attitude, self-efficacy, social support and decrease body weight. Whereas, the use of the Facebook is less time-consuming, more cost-effective, more attractive and provide instantly individual responses. Thus, we recommend using Facebook to aid future health promotion programs.

Funding: Providence University

P160 The Relationship Between Dietary-related Behaviors and Psychological Constructs and Obesity in a Multiethnic Sample

Hiroko Yoshida, PhD, yoshida@nakamura-u.ac.jp, Nakamura Gakuen University Junior College, 5-7-1 Befu, Jonan-ku, Fukuoka, Fukuoka, 814-0198, Japan; J. Maddock, PhD, University of Hawaii

Objective: Differences in dietary intake, obesity and life-style related diseases have been shown in multiethnic cohorts. However, there is little data on Asian/Pacific Islanders' nutrition-related psychological constructs and obesity. We assessed four major ethnic groups in Hawaii: whites, native Hawaiians, Filipinos, and Japanese, on how their dietary-related behaviors and psychological constructs are associated with obesity.

Study Design, Setting, Participants, and Intervention: Data were obtained by random-digit-dial telephone surveys of 2,795 adults ages 18-55 living in Hawaii from one of the four main ethnic groups. Participants were asked questions about their dietary behaviors (fiber, refined grains, fruit and vegetable consumption) and items based on the Theory of Planned Behavior for fruit and vegetables consumption including attitudes, perceived behavioral control and self-efficacy. We examined how these variables were associated with obesity across ethnic groups.

Outcome, Measures and Analysis: We compared differences for BMI and health behaviors by Turkey multiple comparisons, and conducted multivariable logistic regression models.

Results: BMI among native Hawaiians was significantly higher than other ethnic groups. Whites had lower obesity rates than in other groups and scored higher on psychosocial scales. All of the psychosocial constructs had low levels of endorsement in Filipinos and Japanese. Perceived behavioral control in Filipinos were significantly lower than other ethnic groups. Self-efficacy in native Hawaiians was higher than Filipinos and Japanese. Logistic regression results showed that dietary behaviors independently lowered the obesity odds ratio. However, for men, logistic regressions showed different psychosocial predictors for each ethnic group.

Conclusions and Implications: We examined the association between health behaviors and obesity in a multi-ethnic sample in Hawaii. Endorsement of psychosocial variables and predictors of obesity differed across ethnic groups.

Funding: Hawaii State Department of Health, Healthy Hawaii Initiative

P161 Health Risk Factors and Behaviors of Limited-Income Hispanic Women in EFNEP and WIC

Traci Armstrong Florian, MS, RD, tarmstro@cals.arizona.edu, The University of Arizona Cooperative Extension Maricopa County, 4341 East Broadway Road, Phoenix, AZ 85040; D. M. Winham, DrPH, MCHES, Iowa State University

Continued on page S171