P112 (continued)

Design, Setting and Participants: A descriptive survey design among 100 senior secondary two (SS 2) students who were randomly selected from five secondary schools located in education district III of Lagos state, Nigeria.

Outcome Measures and Analysis: Data was collected by means of a pre-tested questionnaire. Academic scores in English language and Mathematics were also collected. Simple percentage, chi square and correlation were used to analyze data.

Results: Findings revealed that majority (74%) of the adolescent students were between 14-16 years, while majority (78%) skips breakfast. Also majority (55%) of the students can participate for only 10 minutes out of a 40-minutes classroom lesson. The results also showed a significant relationship ($r=.456$) between breakfast skipping and academic performance in key subjects (English language and Mathematics) at 0.01 level. A significant relationship also exists between breakfast consumption, classroom participation and academic performance at 0.05 level of significance. Students who do not skip breakfast participate more effectively in class and this has a positive influence on their academic performance.

Conclusions and Implications: Skipping of breakfast is associated with lower nutritional status which consequently leads to low academic performance. Nutrition Education package for adolescents in Nigeria should be developed. Nutritionists can help by communicating correct nutrition information to adolescents through the state schools' board. The schools' canteen should be encouraged to provide low-cost breakfast dishes.

Funding: None.

P113 Analysis of New Child and Adult Care Food Program (CACFP) Cycle Menus by Family Child Care Providers

Mackenzie Ferrante, mackenzie.ferrante@gmail.com, University of Idaho, 307 S. Blaine Street #202, Moscow, ID 83843; M. Sigman-Grant, PhD, RD, University of Nevada; S. Ramsay, PhD, RDN, LD, University of Idaho

Objective: To examine family child care providers' responses to new CACFP cycle menus that included the new USDA Recipes for Healthy Kids.

Design, Setting and Participants: Two groups of family child care providers attending a biannual Food For Kids, Inc. workshop in Reno (n=38) and Las Vegas (n= 39) were recruited. Participants were asked to rank updated CACFP cycle menus on their likelihood of use (following a Likert scale from very likely-very unlikely) and to identify perceived barriers from seven responses (cost, preparation time, ease of preparation, the perceived children's liking, food availability, equipment needed, and storage). Both groups ranked the breakfast and snack menus individually, then were put into groups of two to eight and asked to rank the lunches.

Outcome Measures and Analysis: Descriptive statistics were used to identify participants' likelihood to use menus and ranked responses.

Results: The majority (76%, 19/25) of the lunch menus were likely to be used, and those most likely to be used were simple to prepare. The menus participants identified as least likely to be used were those using USDA Healthy Kids recipes. From the lunch menus identified as least likely to be used, the barriers were preparation time and concern with perceived children's liking. The individually ranked breakfast and snack menus indicated a bimodal distribution between very likely and unlikely to use.

Conclusions and Implications: The updated menus were well received by the majority of child care providers, but menus including USDA recipes were not. Reviewing provider perceptions of USDA recipes and addressing child care providers' reported barriers to use of menus is needed.

Funding: None.

P114 Food Pantries' Response to the Choosing Healthy Options Program: A Mixed-Methods Study in Georgia

Caroline Cahill, MPH, crcahill@gmail.com, Rollins School of Public Health, Emory University, 212 Adair Street, Apartment E2, Decatur, GA 30030; A. Webb-Girard, PhD; J. Giddens, MS, RD, LDN, Atlanta Community Food Bank

Objective: First, to quantitatively assess the nutritional quality of Atlanta Community Food Bank's (ACFB) partner agencies' (PA) food orders before and after implementation of the Choosing Healthy Options Program (CHOP). Second, to understand the factors that influence PAs food ordering process to meet client food preferences and needs.

Design, Setting and Participants: CHOP uses a simple algorithm that designates a 1 (choose frequently), 2 (moderately), or 3 (infrequently) to foods based on nutrient profiles. Food orders of 402 PAs in Atlanta and northwest Georgia were analyzed for the six weeks prior to and following CHOP implementation. Additionally, 9 PAs participated in in-depth interviews.

Outcome Measures and Analysis: Pounds of ordered food by CHOP category and subcategories (e.g., meat, canned vegetables) were analyzed using paired t-tests. In-depth interviews were conducted and analyzed to further understand purchasing motivations, impact of CHOP, client feedback mechanisms, and perceptions of clients' needs.

Results: Overall, agencies increased their percentage of CHOP 1 foods by nearly 14% during the study period. Consistent with PA interview data of clients requesting more meat, PAs increased their pounds of CHOP 1 Meat, though PAs also increased their pounds of CHOP 3 Meat. PAs indicated that CHOP was useful for deciding between food items of the same subcategory that had differing CHOP rankings (i.e., CHOP 2 canned vegetable vs. CHOP 3 canned vegetable). Additionally, only three PAs reported gathering client preference information.

Conclusions and Implications: CHOP seems to have a slight positive impact on PA ordering. More communication is needed between client and provider to offer the clients a more autonomous food assistance experience.

Funding: Center for Ethics, Emory University.