NP5 (continued)

Conclusions and Implications: This Strengthening project is planting the seeds for a multi-year, integrated USDA AFRI project that will significantly impact the food and nutrition needs of children living in rural, low-resourced, American Indian communities.

Funding: 2016-69001-24996

NP6 HomeStyles: Shaping Home Environments and Lifestyle Practices to Prevent Childhood Obesity: A Randomized Controlled Trial

Carol Byrd-Bredbenner, PhD, RD, FAND, bredbenner@aesop.rutgers.edu, Rutgers University, 26 Nichol Avenue, New Brunswick, NJ 08901; John Worobey, PhD, FAPA, Rutgers University; Nobuko (Kay) Hongu, PhD, RD, University of Arizona; Gina Hernandez, MA, Prevent Child Abuse-NJ

Objective: To create an effective, sustainable, population-level obesity-prevention intervention that enables and motivates parents of preschoolers to shape their home environment and lifestyle behavioral practices to prevent excessive weight gain in their children and compare its effectiveness to control condition participants.

Description: Intervention development and implementation was guided by Social Cognitive and Adult Learning Theories, community-based participatory research, and motivational interviewing. The randomized controlled intervention trial is delivered by in-home visitors or online to English and Spanish parents of preschoolers. Focus groups with parents guided development of intervention materials. Cognitive testing with parents was used to refine intervention materials.

Evaluation: Cognitive testing of intervention materials indicated high audience acceptance and satisfaction. Training of home visitation staff delivering the intervention in-home was rated highly satisfactory by staff. Instruments for benchmarking participants and assessing intervention outcomes are valid and reliable. ANCOVA indicated that, compared to control group parents, experimental group parents experienced significant improvements in physical activity behaviors, family mealtime behaviors, self-efficacy for childhood obesity-protective behaviors, and value placed on engaging in childhood obesity-protective behaviors. The vast majority of experimental group parents felt HomeStyles was interesting, useful, easy to use, attractive, and helpful to their families.

Conclusions and Implications: Activities of the final year focused on retaining participants and analyzing data. Parent cognitions and behaviors along with home environments and family lifestyle behaviors that promote healthy child growth and development have the potential to prevent excessive weight gain in preschoolers.

Funding: 2011-68001-30170

NP7 Development and Formative Evaluation of the MENTOR Program for Preventing Childhood Obesity in School-Age Children

Catherine Coccia, PhD, RD, ccoccia@fiu.edu, Florida International University, 11200 SW 8th Street, AHC5 316, Miami, FL 33199; Jessica Moya, BS, RN; Kathryn Brogan-Hartlieb, PhD, RD; Gwyn Senokossoff, PhD, Coker University; Adriana Campa, PhD, RD, Florida International University; Fatma G. Huffman, PhD, RD

Objective: This abstract describes the development and formative evaluation of a school-based nutrition and physical activity intervention program. The aim of the Mentoring, Education, and Nutrition Knowledge for Teachers to Reduce Obesity Risk in Children (MENTOR) program is to determine if a student-teacher focused intervention can enhance classroom nutrition and physical activity content, teacher modeling of positive health behaviors and enrich the health content in the classroom environment.

Description: Program development followed community-based participatory approaches and included three phases: a community needs assessment with in-service (SCTs) and pre-service teachers (PCTs), initial program development and community stakeholder interviews.

Evaluation: Based on feedback from the needs assessment, a multi-modality program was developed: one in-person training session, a project website www.mentorprojectfiu.com to host program curriculum and a social media extension. Community stakeholders (n=4) were asked to review the MENTOR project program content, design and format via in-depth interviews. Overall PCTs and SCTs felt the program was needed, liked the level of detail in the materials and were positive about the format.

Conclusion and Implications: Areas for improvement included the amount of materials needed for teachers to implement some of the lessons in the curriculum along with the amount of time required to implement lessons. This program is currently being evaluated using a pre-post intervention design model. Future directions include the extension of program curriculum and the development of a parent component.

Funding: 2015-69001-23241

NP8 A Social Marketing and Environmental Change Intervention: Get Fruved Years 01-03

Sarah Colby, PhD, RD, scolby1@utk.edu, University of Tennessee, 1215 Cumberland Avenue, Knoxville, TN 37996; Anne Mathews, PhD, RD, University of Florida; Melissa Olfert, DPH, RD, West Virginia University; Kendra Kattelmann, PhD, RD, South Dakota State University; Tandalayo Kidd, PhD, RD, LPN, Kansas State University; Adrienne White, PhD, RD, University of Maine; Tonya Horacek, PhD, RD, Syracuse University; Jesse Morrell, PhD, University of New Hampshire; Lisa Franzen-Castle, PhD, RD, University of Nebraska-Lincoln; Geoffrey Greene, PhD, RD, University of Rhode Island; Carol Byrd-Bredbenner, PhD, RD, Rutgers University; Wenjun Zhou, PhD, University of Tennessee; Onikia Brown, PhD, RD, Auburn University; Karla Shelhutt, PhD, RD, University of Florida

Continued on page S113

NP8 (continued)

Objective: To describe Get Fruved year 03 components and outcomes.

Description: Get Fruved is a health promotion intervention utilizing Community Based Participatory Research to develop and test a social marketing and environmental change intervention to prevent unwanted weight gain among older adolescents. The intervention was developed, refined, and pilot/feasibility tested at four intervention colleges with four other colleges as control. One high school (HS) served as a site for initial development and feasibility testing of the program and data collection tools.

Evaluation: In 2015-2016 (college wave I), 578 non-first year students and 1,148 first year college students (FYCS) completed physical assessments and online surveys and 4,595 FYCS completed online-only surveys. In fall 2016, wave II online survey assessments were completed with 2,491 new FYCS and 237 HS students. Weight was the primary outcome, but since the intervention focused only on obesity-prevention, health-related behaviors, numerous health-related behaviors were assessed.

Conclusions and Implications: Two-year post assessments with college wave I, and one-year post assessments with college wave II will be completed in May 2017. As of year 03, twelve new instruments/methods have been developed/validated, three college courses created, seventy-five presentations conducted, and a college toolkit (providing a systematic approach to promoting health for obesity prevention) has been developed for national testing and dissemination. In year 04, intervention refinement and feasibility testing will occur on eight additional HS campuses. In years 04 and 05 the toolkit will be tested for effectiveness using a Randomized Control Trial design on new college and HS campuses.

Funding: 2014-67001-21851

NP9 Using the Child as Change Agent for Preventing Childhood Obesity in Rural Georgia: Outcome Results on BMI for Age and Home Environment

Marsha Davis, PhD, University of Georgia;
Courtney Still, PhD, RD, courtms@uga.edu, 116E Rhodes Hall, Health Sciences Campus, Athens, Georgia 30602;
Andrea Scarrow, MAL, University of Georgia;
Frances McCarty, PhD, George Washington University;
Richard Christiana, PhD, Appalachian State University;
Richard Lewis, PhD, University of Georgia;
Rebecca Mullis, PhD, RD

Objective: This study aims to test effectiveness of a school-based curriculum, utilizing children as change agents, supplemented by family activities, to prevent childhood obesity.

Description: The Action Pack Families study is a group-randomized trial that follows children from

the beginning of 3rd grade to the end of 5th grade. These analyses consider changes in child body mass index for age from baseline (beginning of 3rd grade) to the end of 5th grade and opportunities for healthy eating and physical activity in the child's home environment.

Evaluation: Data were collected at the beginning of 3rd grade (baseline) and the end of 5th grade in 10 schools. Child height and weight were collected at school by the research team; diet, physical activity, and home environment factors were self-reported by parents using a paper-pencil survey. We will present data from baseline to the end of 5th grade, focusing on child weight status.

Conclusions and Implications: Using children as change agents may be an effective way to alter the home environment and prevent childhood obesity.

Funding: 2012-68001-19625

NP10 Engaging Youth as Advocates to Create Healthy Snacking Zones Around Rural Schools - Year 5

Nancy Findholt, PhD, RN, findholt@ohsu.edu, Oregon Health & Science University, One University Boulevard, La Grande, OR 97850; Betty Izumi, PhD, MPH, RD, Portland State University; Jackilen Shannon, PhD, MPH, RD, Oregon Health & Science University; Thuan Nguyen, PhD, MD; Carole Smith, MEd, Oregon State University Extension Service

Objective: This quasi-experimental study evaluated an intervention that engaged 4-H youth participants in advocacy for "healthy snacking zones" within 5 elementary/middle schools and nearby small food stores in a rural Oregon county.

Description: Youth engagement occurred through participation in 4-H "SNACZ clubs," which were established in each intervention school. Youth club members received nutrition lessons and, with support from adult club leaders, planned and implemented outreach projects to promote healthy snacking among their peers and encourage environmental change within their schools and the food stores.

Evaluation: Mixed methods were used to assess changes in school and store environments, as well as changes in children's snacking behaviors and BMI.

Conclusions and Implications: After the SNACZ program, intervention teachers had greater overall nutrition knowledge ($p=0.098$) and were less likely to use food rewards for students, including sweetened drinks (0.053), than teachers in control schools. Mean availability of healthy snacks/beverages in food stores near intervention schools increased by 21%, while mean availability of these products in stores near control schools decreased by 19%. Students' mean consumption of unhealthy snacks/beverages outside of school decreased in both intervention and control sites, but the magnitude of the change was substantially greater in

Continued on page S114