P46 Diabetes Prevention Program Results in Immediate Lifestyle Changes Independent of Weight Loss

Hannah Dentry, BS, mudahogo@gmail.com, University of Maryland, 112 Skinner Building, College Park, MD 20743; Alison Massey, MS, LDN, CDE, University of Maryland; Margaret Udaogora, PhD, MD

Objective: This study measured the immediate impact of lifestyle changes taught by the Diabetes Prevention Program (DPP), understanding that changes may not be reflected in weight loss.

Study Design, Settings, Participants: This study followed 34 adults in the DPP. Participants received classes from the CDC's curriculum and emails with follow-up information and videos, created outside of the CDC's curriculum. Surveys were created to measure the immediate impact of topics covered in each period (2, 3 and 4 months) by comparing participants' wellness practices before and after the intervention.

Measurable Outcome and Analysis: Wellness practice ratings were used to evaluate the immediate impact of the DPP. Areas of interest included compliance with 2015 Dietary Guidelines, physical activity, and stress management. Average weight loss was recorded.

Results: Participants filling half their plates with vegetables increased from 23% to 55%. Participants engaging in 30 minutes of physical activity increased from 18% to 60%. Participants who "Almost Always" ate in response to stress decreased from 33% to 0%. Of the participants who received emails, 80% watched the videos. 67.6% of participants lost weight, ranging from -6.4% to 10.6% weight loss and an average weight loss percent of 0.30%.

Conclusion and Implications: Participants immediately implemented the lifestyle changes they learned from the DPP, although average weight loss was low. To better understand the impact of the DPP, future studies could evaluate the long term maintenance of lifestyle changes and compare lifestyle changes made by those who received the emails and videos, versus those who did not.

Funding: None

P47 Dietary and Exercise Counseling Provided to Clients at Risk for Cardiovascular Disease: Findings from National Health Survey

Oyinlola Babatunde, PhD, MPH, RD, babatundeo@ecu.edu, East Carolina University, Health Science Building, 3310J MailStop #668, Greenville, NC 27834; Charles Ellis, PhD, East Carolina University; Hui Bian, PhD

Objective: To examine the provision of dietary and exercise counseling to clients with comorbid conditions

who are at risk for cardiovascular disease (CVD), and the relationship to BMI.

Study Design, Setting, Participants: Secondary analysis of the 2013 National Ambulatory Medical Care Survey (NAMCS) data for respondents ≥ 30 years and obese (BMI ≥ 30) with diabetes, hypertension and/ hyperlipidemia). Respondents data (n=1574) were compared to examine reports of dietary and exercise counseling provided.

Measurable Outcomes and Analysis: Proportion of dietary and exercise counseling reported by respondents. Univariate comparisons of respondents characteristics were completed using descriptive and chi-square statistics. Binary logistic regression conducted to determine the odds of receiving counseling after controlling for relevant covariates.

Results: The mean age of respondents included in analysis was 61.1 years (9.8 SD) with mean BMI=39.3 (6.8 SD). Only 31% of the respondents received dietary and/ exercise counseling despite their comorbid conditions. Chi-square test was significant for BMI groups, $\chi^2(2, N=1574) = 6.44, p=.04$ and ethnicity, $\chi^2(3, N=1574) = 12.0, p=.01$. Further logistic regression analysis revealed minority groups were more likely to receive counseling (White=reference) -Blacks (OR=1.4, $p=.05$, 95% CI=1.01-1.96) and Hispanics (OR=1.7, $p=.01$, 95% CI=1.15-2.47).

Conclusions and Implications: Findings indicate overall percentage receiving dietary and exercise counseling is very low despite their comorbid conditions. Additionally, minority groups are more likely to receive counseling compared to Whites. In defining the future of care to reduce health disparities and promote healthy aging there is need for further studies to better understand why obesity and comorbid conditions are not decreasing among minority groups despite this effort.

Funding: None

P48 Differences in Dietary Behaviors and Nutritional Risk by Living Status in Korea Adults

So-Young Nam, PhD, Inha University; Soo-Kyung Lee, PhD, RD, skleenutrition@inha.ac.kr, Inha University, 100 Inharo, Incheon, Namgu, South Korea 22212

Objective: People living alone have been reported to eat differently from those living with others. Rapid increase in the number of people living alone has become a social issue in South Korea, yet few studies have examined the issue. This study aimed to examine whether and how living status was associated with dietary behaviors and nutritional risk among Korean adults.

Study Design, Setting, Participants: A cross-sectional study was conducted using the Korean National Health and Nutrition Examination Survey (KNHANES 2005~2013)

Continued on page S42