P11 (continued)

Results: The dietary recommendations and goals most frequently mentioned by participants were limiting sodium and increasing intake of fruits and vegetables. Many participants reported that high food prices and a limited budget made it difficult to follow their diet and make healthy choices when grocery shopping. The food environment, particularly the high-salt content in food products and the promotion of unhealthy food in stores, presented additional challenges. Participants suggested a variety of ways to make following their diet, shopping, and cooking easier including online grocery ordering and delivery, pre-prepared ingredients, and assistance with meal planning, shopping, cooking, and cleaning up. The idea of a medically tailored home-delivered meal service was well-received by most study participants.

Conclusions: Our findings provide a starting point for developing programs and policies to help older, lower SES HF patients follow their diet and improve their health, thereby reducing health disparities. Since finances are a key concern, programs and policies need to be designed with cost in mind.

Funding: J.B. Hawley Research Award from the Division of Epidemiology and Community Health at the University of Minnesota

P12 Cooking Self-Efficacy, Perceived Health Status, and Fresh Produce Purchasing Criteria Among Faculty and Staff

Georgianna Mann, PhD, grmann@olemiss.edu, University of Mississippi, PO Box 1848, Sorority Row, Lenior 220, University, MS, 38677; Laurel Lambert, PhD, LD, RDN, University of Mississippi; Kritika Gupta, MS, University of Mississippi; Selby Greer, MS, University of Mississippi

Background: National health guidelines emphasize the nutritional benefits of fruit and vegetable (FV) consumption. Positive associations between FV consumption, health awareness, and cooking skills have been noted in previous work. However, little is known about this relationship among university faculty and staff.

Objective: To explore the relationship between cooking skills, health perceptions, importance of FV consumption, and FV purchase criteria among university faculty and staff by gender.

Study Design, Setting, Participants: A random representative sample of faculty and staff at a large Southern university were recruited to complete an online survey in spring 2018 (n = 175).

Measurable Outcome/Analysis: Questions included cooking skills, perceived health status, benefits of produce consumption, produce purchasing criteria, and demographic data. Associations between cooking skills, self-health assessment, and fresh produce buying preferences were evaluated using Pearson's correlation test. Independent t-tests were used to determine differences by gender. Significance was determined by $P \leq .05$.

Results: Health perceptions were positively correlated with perceived health benefits of fresh produce ($P < .05$),

non-GMO/Organic ($P < .05$), and locally grown ($P < .05$). Female participants reported higher cooking skill levels. Cooking skills were positively associated with grocery shopping responsibilities ($P < .01$), shopping at the farmer's market ($P < .01$), and home gardens ($P < .05$).

Conclusions: In this study, females responded with the highest cooking skill levels which positively correlated with importance of fresh produce, specific purchasing preferences, and self-perceived health status. While overall cooking skill level in this sample was high, it is likely that those interested in healthy eating were more likely to complete the survey. Universities could play an important role in improving fresh produce consumption among faculty and staff by providing support and resources in the form of cooking classes through university programs such as nutrition departments, university wellness programs, and community venues.

Funding: None.

P13 Social Network Analysis of Urban Church Food Pantries

Tracy Noerper, PhD, LDN, RDN, Tracy.Noerper@lipscomb.edu, Lipscomb University, One University Park Dr, Nashville, TN, 37204; Janet Colson, PhD, RDN; Andrew Owusu, PhD

Background: Examining social networks helps to explain the connectedness of individuals, groups or communities, and is useful in demonstrating the potential that lies within and between networks. When evaluating social networks, relationships are viewed as "nodes" (eg individuals or points) and their connecting "ties" (eg the relationships between the nodes). Detailing food pantry and client "connectedness" through social network analysis can assist researchers and policymakers to enhance efficiency and effectiveness of the informal emergency food system.

Objective: The objective of the study was to examine the degree to which urban church food pantries are connected to the clients they serve using the social network theory.

Study Design, Setting, Participants: The research design was a population-based, cross-sectional study using a snowball sampling method of church food pantries in Nashville, Tennessee, from 2015–2018. The study population included 93 urban churches with 46 having on-site food pantries.

Measurable Outcome/Analysis: Zip codes for church-based pantries and clients served were collected via an electronic survey tool sent to urban churches. Social network ties between church food pantries and clients were analyzed utilizing UCINET 6 software.

Results: From the social network analysis a sociogram (eg visual matrix) was developed depicting 56 nodes (26 churches; 30 client zip codes) and 148 ties. Five large churches (membership ≥ 300), 13 medium-sized churches (membership = 100 to 299), and 8 small churches (membership ≤ 100) were identified within 23 zip codes.

Continued on page S22

P13 (continued)

Large churches had 31 ties, medium-sized churches had 98 ties and small churches had 19 ties. Two central, medium-sized churches had the most ties with 20 and 30, respectively.

Conclusions: Although it was assumed that larger churches would have more social network ties, church size was not a limiting factor in serving urban areas through food pantries. Further research is needed to evaluate if current church food pantry social networks ensure vulnerable populations are being adequately served.

Funding: None.

Behavior and Education Theory

P14 Mindful Eating Scores Differ by Body Mass Index and Self-Reported Health Status Among College Students

Opeyemi Adewumi, BS, oadewumi@crimson.ua.edu, University of Alabama, 404 Russell Hall, Box 870311, Tuscaloosa, AL, 35475; Caroline Brantley, MS, LDN, RDN, University of Alabama; Linda Knol, PhD, LDN, RDN, University of Alabama

Background: The prevalence of overweight and obesity continues to rise among college students. College students are also adopting mindfulness habits, such as yoga and mindful meditation, at higher rates than the general population. Mindful eating is a “non-dieting” approach that focuses on paying attention to the body’s internal hunger and satiety cues, while enjoying the food without distraction.

Objective: The purpose of this study was to understand whether mindful eating is related to weight and health status among college students.

Study Design, Setting, Participants: Six hundred and ninety college students participated in an online survey.

Outcome Measures/Analysis: The survey included: the Mindful Eating Questionnaire (MEQ), a validated global health question, self-report of weight and height, and demographics. Mindful eating was reported on a five-point Likert scale where higher scores represented higher mindful eating practices. Multiple regression analysis was used to test for significant differences in mindful eating scores by weight status and health status after adjusting for demographics.

Results: Among respondents, 19.6% rated their health as fair or poor and 67.8%, 20.7%, and 11.5% were classified as healthy weight, overweight, or obese, respectively. After adjusting for demographics, MEQ scores were significantly higher among students reporting their health as excellent/very good/good vs fair/poor (3.4 ± 0.4 vs 3.2 ± 0.4 , $P < .0001$). MEQ scores were significantly different between weight status categories where students in the healthy weight range (3.4 ± 0.4) had significantly higher scores than their overweight (3.3 ± 0.1 , $P = .003$) or obese counterparts (3.2 ± 0.4 , $P < .0001$).

Conclusions: Among college students, mindful eating scores were slightly higher for students at a healthy weight and reporting good health compared to their counterparts. Thus, mindful eating interventions may be an effective strategy for addressing overweight and obesity in a university setting.

Funding: None.

P15 Self-Efficacy and Psychosocial Determinants of Obesity Prevention Behaviors in Caucasian Americans

Doreen Liou, EdD, RDN, lioud@montclair.edu, Montclair State University, 1 Normal Ave, Montclair, NJ, 07043; Lauren Kulik, BA, Montclair State University

Background: The obesity epidemic is a salient health issue affecting Caucasian Americans in the United States. Food consumption patterns, dietary choices, and physical activity behaviors influence weight-related outcomes in individuals. Psychosocial theories provide a foundation in understanding mediators of health behaviors.

Objective: To identify psychosocial determinants of obesity prevention behaviors in Caucasian Americans residing in New Jersey.

Study Design, Setting, Participants: A cross-sectional survey design was used in which a convenience sample of 174 participants (18 to 40 years old) completed a validated online, self-administered questionnaire. Male and female participants were recruited from universities, religious organizations, and cultural institutions, representing a wide range of educational and socioeconomic backgrounds in New Jersey.

Measurable Outcome and Analysis: Nineteen behaviors related to obesity risk reduction were measured along with self-efficacy and psychosocial constructs derived from the Theory of Planned Behavior. Participants indicated their adoption of behaviors over the previous month reflecting food practices, portion size control, consumption of plant-based foods (eg fruits, vegetables, whole grains), and engagement in physical activity. Descriptive statistics and regression analyses were conducted for the entire sample. T-tests were applied for individuals categorized as high vs low self-efficacy groups.

Results: The mean age of the study participants (70% female) was 26.4 (SD = 7.0) years with an average body mass index of 24.7 (SD = 4.8). Regression analysis showed that 37.5% of the variance in obesity risk reduction behaviors was accounted solely by self-efficacy. T-tests indicated that there was a higher frequency of adoption of 17 health behaviors within the high self-efficacy group as compared to their low self-efficacy counterparts. Notable mean differences in behavior included eating home-cooked meals ($P < .05$), limiting portion sizes of food ($P < .001$), eating fruits and vegetables ($P < .001$), engaging in physical activity ($P < .001$), and monitoring body weight ($P < .01$).

Continued on page S23