P90 (continued)

whether use of each category of food label changed over time.

Results: The use of all 4 label categories significantly increased from 2017 to 2019, while the proportion of respondents stating that they did not read food labels decreased. Reading of ingredient lists (86%) and nutrition panels (85%) were more frequently reported compared to front of package (74%) and production practice (72%) label claims.

Conclusions: Food label usage is increasing in general, and consumers are increasingly reading both front of package claims (organic, natural, low sodium) and back of package information (ingredients, nutrition facts panel). As consumer demand for “clean label” products increases, more research is needed to better understand the health impacts of such products and the best methods for labeling them.

Funding: None.

Basic Food and Nutrition Knowledge

P91 Pilot Test of Supporting Wellness at Pantries (SWAP): Clients Chose Healthier Foods after Implementation

Abiodun T. Atoloye, PhD, cPHN, abiodun.atoloye@uconn.edu, Rudd Center for Food Policy and Obesity, University of Connecticut, One Constitution Plaza, Hartford, CT 06103; Sarah L. McKee, MS, University of Connecticut, Department of Human Development and Family Sciences; Eminent Abebe Gurganus, MS, University of Connecticut, Department of Human Development and Family Sciences; Ran Xu, PhD, University of Connecticut, Department of Allied Health Sciences; Katie Martin, PhD, Foodshare Institute for Hunger Research and Solutions; Marlene B. Schwartz, PhD, Rudd Center for Food Policy and Obesity, University of Connecticut

Background: The US food banking network serves about 46.5 million individuals with food insecurity yearly. People living with food insecurity are at risk of diet-related chronic diseases and evidence suggests that foods available in food pantries are mostly nutritionally poor. In response, the Supporting Wellness at Pantries (SWAP) system ranks food into green, yellow, and red categories based on the levels of saturated fat, sodium, and sugar. SWAP is designed to increase the supply and demand for healthy food in pantries.

Objective: To assess the nutritional quality of clients' selections before and after implementing the SWAP system.

Study Design, Setting, Participants: Availability of green, yellow, and red items was captured pre- and post-intervention using a pantry inventory assessment. For the intervention, foods in the pantry were rearranged by SWAP ranking and signage was added to educate clients about which items to choose often (green), sometimes (yellow), or rarely (red). Two-hundred twenty-two clients participated in the pre-post study.

Measurable Outcome/Analysis: Clients' selections were recorded and the proportion of green and red

selections in total and per food group were calculated. Primary analyses compared the proportion of green and red foods chosen by clients before and after SWAP implementation using independent t-tests. A regression model was used to compare clients' selection pre to post while controlling for food pantry inventory.

Results: There was not a significant change in the overall pantry inventory over time. However, the proportion of healthy (ie, green) food selections significantly increased and less healthy (ie, red) food selections significantly decreased. The overall improvement in nutritional quality of the clients' carts remained significant while controlling for the pantry inventory.

Conclusions: Results suggest that nutrition information provided through stoplight signage and clear messages on items to choose often, sometimes, and rarely can shift client choices in the pantry setting.

Funding: Institute for Collaboration on Health, Intervention, and Policy (InCHIP) Seed Grant.

Agricultural Production and Food Systems

P92 Controlling for Confounding Factors in a Multi-Component Farmers' Market Intervention

Hayley Heino, MS, California State University, Chico's Center for Healthy Communities; Jenni Dye, MS, RDN, California State University, Chico's Center for Healthy Communities; Naomi Stamper, MS, California State University, Chico's Center for Healthy Communities; Robin Donatello, PhD, California State University, Chico's Data Science Initiative; Stephanie Bianco, MS, RD, sbianco@csuchico.edu, California State University, Chico's Center for Healthy Communities, 25 Main St, Suite 101, Chico, CA, 95928-5388

Background: Despite federal funding to encourage SNAP EBT use at farmers' markets, only \$22 million or 0.03% of the \$70 billion total SNAP dollars are spent in these direct-to-consumer channels.

Objective: The objective of this study was to determine if confounding factors such as weather or other market activities have an impact on SNAP sales compared to typical market days.

Study Design, Setting, Participants: This 4-year, quasi-experimental study used a multi-component intervention at a medium-sized, weekday, year-round farmers' market in northern California. Total SNAP sales were collected over 54 baseline and 136 intervention market days, respectively. SNAP recipients were the target population, however several intervention components were available to all market shoppers. The multi-component intervention included SNAP outreach (ie, on-site eligibility screening, SNAP application assistance, and SNAP material distribution), SNAP signage, financial incentives (ie, Market Match), and nutrition education including food tastings highlighting market produce, recipe cards, posters, and market education.

Continued on page S60