O12 (continued)

how peer educators perceive program recipes and related participants' recipe acceptability and preparation behaviors.

Objective: To examine peer educators' acceptability of recipes through evaluation of sensory attributes (appearance, flavor, texture, and saltiness or sweetness), preparation behaviors, and participants' perceived behaviors.

Study Design, Setting, Participants: Eight online surveys were conducted with peer educators (100% female, 47% African American, 14-28 responses per recipe) in the University of Georgia EFNEP and SNAP-Ed programs. Educators completed the surveys independently in a location of their choice. Survey links were sent to educators in a single document, and weekly reminders paced them to complete 2 surveys per week.

Measurable Outcome/Analysis: Sensory acceptability of 16 recipes was evaluated on a 9-point hedonic scale (0=Dislike Extremely; 5=Neither Like nor Dislike; 9=Like Extremely). Preparation intent and behaviors of educators and participants were evaluated on a similar scale. Standard range of acceptability is ≥ 7 .

Results: Thirteen recipes received acceptable ratings on most sensory attributes and preparation questions. Three recipes (Curly Noodles Supreme, Easy Cheesy Broccoli Soup, and Ranch Sauce) received unacceptable ratings. Across all recipes, recipe acceptability, preparation intent, and program participant perception were not significantly different by educators' program, length of employment, and quadrant district in Georgia.

Conclusion: The findings suggest that peer educators for the nutrition education of low-income Georgians reported overall acceptable sensory attributes and preparation behaviors related to the program recipes and indicate areas for improvement in select recipes. Sensory evaluation methodology could be used to improve nutrition education recipes for higher acceptability and healthier recipe preparation behaviors in low-income participants.

Funding: Supplemental Nutrition Assistance Program - Education.

School Age Children/School-Based Programs

O13 Acceptability, Perceived Benefits, and Unintended Consequences of a Virtual Nutrition Intervention for Adolescents with Autism Spectrum Disorder (ASD)

Acadia Buro, MS, acadia@usf.edu, University of South Florida, 13201 Bruce B Downs Blvd, Tampa, FL, 33612; LaShae Rolle, BS, University of South Florida; Heewon Gray, PhD, RDN, University of South Florida

Background: Adolescents with ASD exhibit an increased prevalence of problematic eating behaviors, such as food selectivity, or consuming a narrow range of foods, and consume more energy-dense foods and fewer fruits and vegetables than typically developing children.

Objective: The purpose of this study was to examine the acceptability, perceived benefits, and unintended

consequences of a virtual implementation of BALANCE (Bringing Adolescent Learners with Autism Nutrition and Culinary Education), an 8-week theory-driven nutrition intervention for adolescents with ASD.

Study Design, Setting, Participants: A qualitative study design was used. Twenty-seven adolescents aged 12-20 years who attended the BALANCE intervention and their parents participated in the study. Twenty-one parents participated in an interview. Six focus groups with 1 to 5 adolescents in each group were conducted. Data were collected virtually via Microsoft Teams.

Measurable Outcome/Analysis: Focus groups and interviews were audio-recorded and transcribed. Data were analyzed for a priori and emergent themes regarding intervention acceptability (likes, dislikes, satisfaction, and suggestions for improvement), perceived benefits, and unintended consequences.

Results: The intervention was generally acceptable to participants. Parents described, "I was so impressed," "I feel like [he] was so happy," and "he is more open to try new things." Themes for acceptability included "virtual format," "group setting," "autonomy/independence," "sensory components," "interaction," "reinforcement," and "parent component." Themes for perceived benefits included "diet changes," "healthy weight," "knowledge/awareness," "behavioral skills," "self-efficacy," "outcome expectations," "outcome expectancies," and "other lifestyle changes." "Anxiety/discomfort" during intervention lessons was an emergent theme regarding unintended consequences.

Conclusion: Our findings suggest that a virtual implementation of BALANCE was acceptable according to adolescents with ASD and their parents based on their reported perceptions and feelings about the intervention. Based on participant feedback, many adolescents with ASD may benefit from small group interventions, and virtual interventions are a convenient option for some adolescents and parents.

Funding: University of South Florida.

O14 Sugar-Sweetened Beverage Consumption Among Youth in North Carolina

Lindsey Haynes-Maslow, PhD, MHA, lhaynes-maslow@ncsu.edu, North Carolina State University, 512 Brickhaven Dr, Room 240c, Raleigh, NC, 27699; Jonathan Blitstein, PhD, RTI International; Sarah Ray, MA, RTI International; Jenna Brophy, MPH, RTI International; Michael Hayes, BA, RTI International; Micaela Brewington, BS, RTI International

Background: Sugar-sweetened beverage (SSB) consumption among youth contributes to diet-related chronic disease including obesity, type 2 diabetes, and poor oral health.

Objective: Collect data to inform the development of a social marketing campaign aimed at reducing SSB consumption among youth age 11-17 in North Carolina (NC).

Study Design, Setting, Participants: Cross-sectional survey data collected from September 9, 2020 through

Continued on page S7