P28 (continued)

Conclusion: The researchers, despite being experienced nutrition educators, were surprised at the number of unfamiliar H&S, and that many were used for medicinal purposes yet were grouped with H&S used for culinary purposes, sometimes even on the same shelf. These findings support the following implications: 1) inclusion of community members in nutrition education materials development is essential for cultural relevance; 2) nutrition educators could benefit from additional cultural cuisine training; and 3) approaching the study of non-Western cultures with an ethnocentric bias may prohibit a thorough understanding of how people may use H&S at home.

Funding: None.

P29 Interest and Acceptability of a Clinic Based EFNEP Intervention in English and Spanish Speaking Parents for Pediatric Obesity Prevention

Mical Shilts, PhD, MS, shiltsm@csus.edu, California State University, Sacramento, 6000 Jst, Sacramento, CA, 95819-6053; Karina Diaz Rios, PhD, RDN, University of California, Merced; Lenna Ontai, PhD, University of California, Davis; Katherine Panarella, MPH, MS, University of California, Davis, Agriculture and Natural Resources; Christiana Drake, PhD, University of California, Davis; Dennis Styne, MD, University of California, Davis Children's Hospital; Samantha Navarro, BA, University of California, Merced; Adrian Loera, BS, University of California, Merced; Louise Lanoue, PhD, University of California, Davis; Marilyn Townsend, PhD, RD, University of California, Davis

Objective: Obesity continues to disproportionately impact low-income, ethnically diverse youth. Existing government-funded nutrition education programs and pediatric medical clinics can collaborate to target and deliver relevant interventions. The objective was to explore interest and acceptability of community nutrition intervention collocated in medical clinics.

Use of Theory: Social Cognitive (SCT) and goal setting theories.

Target Audience: Low-income English and Spanish-speaking parents with young children.

Program Description: Physicians referred parents of pediatric patients to the intervention. The intervention consisted of 8-weekly, 1.5 hour nutrition, activity and food-related parenting content anchored with guided goal setting, SCT constructs, and story telling. Existing Expanded Food and Nutrition Education (EFNEP) educators delivered the intervention in medical clinics.

Evaluation Methods: To assess interest and acceptability, referral, enrollment, and attendance data were collected and parent focus group interviews were conducted. Descriptive statistics and chi-square analysis were conducted on the quantitative data and interviews were recorded, transcribed and organized for content analysis.

Results: Twenty intervention series were conducted in English (n = 12) and Spanish (n = 8) at 5 medical clinics by 4 EFNEP educators. More than 1/3 of the parents (37%; n = 709) expressed interest in the intervention after physician referral and 74% (n = 195) enrolled in the intervention. A greater proportion of referred Spanish-speaking parents expressed interest ($P < 0.001$) and completed 4 or more intervention sessions ($P < 0.0001$) compared to English-speaking parents. A total of 26 focus group interviews (n = 65) were conducted with the 97% of the parents said they enjoyed participating in the intervention. The intervention component most commonly mentioned as useful was guided goal setting.

Conclusion: Results from this study indicated that parents referred to the intervention were interested and attended with greater appeal to Spanish-speaking parents. Acceptability of the intervention can be attributed to the tailoring of the content and delivery by culturally competent educators.

Funding: NIFA.

P30 Interface of Neighborhood Residence on Obesity Prevention Behaviors and Self-Efficacy in Chinese Americans

Doreen Liou, EdD, RDN, lioud@montclair.edu, Montclair State University, 1 Normal Ave, Montclair, NJ, 07043; Jessica Karasik, BA, Montclair State University

Background: The obesity epidemic is a prominent health issue affecting Chinese Americans in the United States. An individual's living environment affects dietary choices and consumption patterns, impacting weight-related outcomes. The important construct of self-efficacy provides a foundation for nutrition educators in understanding mediators of health behaviors.

Objective: To identify if self-reported neighborhood of residence ('high income' vs 'middle-low income') elucidate differences in obesity prevention behaviors and self-efficacy in Chinese Americans residing in Los Angeles county and the New York metropolitan area.

Study Design, Settings, Participants: A cross-sectional survey design was used where a convenience sample of 650 participants (18-60 years old) completed a validated, self-administered questionnaire. Participants were recruited from universities, cultural institutions, and churches, representing a wide range of backgrounds.

Measurable Outcome and Analysis: Nineteen behaviors related to obesity risk reduction were measured with 9 items measuring self-efficacy for these behaviors. Participants indicated adoption of behaviors over the month reflecting food practices, portion size control, and consumption of plant-based foods. Descriptive statistics were conducted for the entire sample while *t* tests were applied for categorized neighborhoods ('high income' vs 'middle-low income').

Results: The mean age of the study participants (67%

Continued on page S38

P30 (continued)

female) was 36.3 (SD = 14.5) years. Higher frequency of adoption of 6 health behaviors within the 'high income group' as compared to counterparts was indicated by *t* tests. Notable mean differences in behavior included choosing steamed over fried foods ($P < 0.01$), using small amounts of oils ($P < 0.05$), eating healthful snacks ($P < 0.05$), and following traditional healthful Chinese food patterns ($P < 0.01$). Stronger self-efficacy levels in the 'high income group' for 7 out of 9 items reflecting the performance of dietary behaviors were also indicated by *t* tests.

Conclusion: Nutrition professionals must assess client's living environments in the adoption of obesity prevention behaviors. In fostering behavioral confidence, investigating the impact of individuals' neighborhood in various residential areas are warranted.

Funding: None.

P31 Management of Disordered Eating Behavior Risk in Clients Varies Among Outpatient Dietitians: A Qualitative Thematic Analysis

Dagny Larson, BS, RD, LD, dagnynlarson@austin.utexas.edu, Department of Nutritional Sciences, The University of Texas at Austin, 200 W. 24th St, Gearing Hall, Room 313, Mail Stop A703, Austin, TX, 78712; Jacqueline Henning, BS, Department of Nutritional Sciences, The University of Texas at Austin; Marissa Burgermaster, PhD, Department of Nutritional Sciences and Dell Medical School, The University of Texas at Austin

Background: Sixty-two percent of dietitians recommend nutrition apps to patients, with calorie-tracking apps the most popular. While previous research has indicated common characteristics of nutrition app users, little is understood about how client characteristics impact dietitians' decisions about recommending apps. Addressing this relationship would provide guidance for dietitians to make evidence-based practice decisions regarding which clients would benefit.

Objective: To explore the role of client characteristics in outpatient dietitians' decisions to recommend apps.

Study Design, Setting, Participants: Twenty- to 60-minute semi-structured interviews conducted by phone, videoconferencing, or in-person from November 2019 to July 2020 with U.S. outpatient dietitians ($n = 20$).

Measurable Outcome/Analysis: Interviews investigated dietitian experiences surrounding their app decision-making process, including the client-, app-, and practice-related factors involved. Two coders conducted a qualitative thematic analysis of the client-related factors.

Results: Dietitians years in practice ranged from 2 to 42 year and they identified weight management and diabetes as top counseling topics. We identified 3 client-related themes: motivation, tech/health literacy, and disordered eating behaviors. While clients' motivation and literacy level may impact the dietitian's decision, the presence of disordered eating behaviors more consistently impacted these decisions, leading dietitians to avoid or carefully

select apps for clients, especially calorie-tracking apps. Many participants perceived that apps would benefit or harm clients based on a nuanced set of characteristics, such as being numbers-oriented vs obsessive. Several participants reported they avoided the use of calorie-tracking apps for all clients, including those without concerning characteristics or behaviors, due to worries that their use would increase risk of developing disordered eating behaviors.

Conclusion: While several client features influenced dietitians' app decisions, disordered eating behaviors appeared to be the most consistent characteristic leading to avoidance or adjusted use. Despite this consistency, dietitians differed regarding whether apps promote these behaviors in all clients. Further research to understand this relationship is needed to guide dietitians and app design.

Funding: None.

P32 Self-Efficacy Among Students in Various Dietetics Programs

Kristen Heitman, MS, RD, LD, The Ohio State University; Christopher Taylor, PhD, LD, RDN, FAND, taylor.1043@osu.edu, The Ohio State University, 1675 Demaret Ln, Columbus, OH, 43228

Background: In educational programs, student perceptions are often overlooked; yet, have been shown to contribute to motivation, engagement, and academic performance. Conversely, a lack of self-efficacy can be a barrier to professional and effective clinical practice.

Objective: The objective of this research is to explore student self-efficacy in various dietetics programs.

Study Design, Setting, Participants: A cross-sectional study design assessed self-efficacy among dietetics students ($n = 44$) in 4 programs: coordinated undergraduate, coordinated graduate, dietetic internship, and the Future Education Model Graduate Program. Students were classified as first year of the program or subsequent year, as well as first year or subsequent year(s) as undergraduate and graduate students.

Measurable Outcomes/Analysis: The Self-Efficacy Scale for Nutrition Students in the Clinical Setting is a survey consisting of 20 statements based on ACEND competencies for programs. Survey statements are rated on a 4-item Likert scale ranging from 'not at all confident' to 'fully confident.' This survey was administered electronically and mean self-efficacy scores across all items were compared using independent samples *t* test and ANCOVA.

Results: Students in the first year of a dietetics program had significantly lower scores (2.9 ± 0.53) than in students in subsequent years (3.4 ± 0.41 , $P = 0.004$). First year graduate students had significantly lower scores (2.79 ± 0.38) than second year graduate (3.37 ± 0.46) and undergraduate students (3.47 ± 0.29 , $P = 0.008$). Areas of highest confidence among all students were ability to abide by the code of ethics, collect food and nutrition-related history,

Continued on page S39